Psychreg Journal of Psychology OPEN ACCESS

Volume 10, Issue 1, June 2026 ISSN 2515-138X | pjp.psychreg.org

REVIEW ARTICLE

Mindfulness-based interventions and youth mental health: A review of recent evidence and future directions

Jailey Hall

University of Pikeville (United States)

Correspondence: jaileyedyn@hotmail.com

Cite as: Hall, J. (2026). Mindfulness-based interventions and youth mental health: A review of recent evidence and future directions. *Psychreg Journal of Psychology*, *10*(1), 1 to 7. https://doi.org/rm2g

ABSTRACT

Mindfulness has emerged as an influential framework for enhancing mental health, with robust evidence among adults and growing interest in its application to adolescent populations. This perspective highlights the potential of mindfulness-based interventions (MBIs) as a practical and accessible response to the global adolescent mental health crisis. Rather than conducting a systematic review, we draw on selected empirical and theoretical contributions to argue that mindfulness can equip adolescents with essential coping skills during a critical period of psychological development. Adult research has demonstrated consistent effects of MBIs, including mindfulness-based stress reduction (MBSR) and mindfulness-based cognitive therapy (MBCT), on outcomes such as stress, depression, anxiety, and overall well-being. Emerging evidence with adolescent samples points to comparable benefits, including improvements in emotional regulation, resilience, and interpersonal functioning. MBIs may hold particular value for underserved populations, where structural and socioeconomic barriers to traditional mental health care remain pronounced. Cultural relevance, accessibility, and sustained engagement are each identified as important considerations when adapting these interventions for diverse youth contexts. This perspective argues that integrating MBIs into schools, community programmes, and digital platforms represents a promising pathway for addressing unmet mental health needs among young people. Future efforts should prioritise cultural adaptation, flexible delivery formats, and strategies to support long-term engagement. By drawing on insights from both adult and adolescent research, we position mindfulness not solely as a therapeutic tool but as a preventive and developmental resource with broader public health implications for youth.

Keywords: adolescent mental health; mindfulness-based interventions; psychological well-being; resilience; youth coping

Mindfulness has become a central theme in contemporary psychology, with decades of research demonstrating its potential to reduce stress, regulate emotions, and promote well-being (Ogwuche et al., 2020). Initially established through interventions such as mindfulness-based stress reduction (MBSR; Kabat-Zinn, 1994) and mindfulness-based cognitive therapy (MBCT; Segal et al., 2002), mindfulness has moved well beyond clinical settings to influence education, healthcare, and workplace practices.

Much of this work has centred on adults. A pressing question remains: how can mindfulness be meaningfully applied to adolescents, a group navigating significant developmental challenges and mounting mental health risks? Adolescence is marked by identity formation, social complexity, and heightened vulnerability to anxiety and depression (World Health Organization, 2021). Traditional mental health services often struggle to reach young people effectively, particularly in underserved contexts where need is frequently greatest (Bear et al., 2024).

This perspective argues that mindfulness offers a practical and adaptable response to this gap. By cultivating present-moment awareness and emotional resilience, MBIs can equip adolescents with coping skills that extend beyond symptom reduction. Rather than providing a comprehensive review, this paper draws on selected studies to highlight both the promise and the limitations of adolescent MBIs, and proposes directions for integrating mindfulness into schools, community programmes, and digital platforms with attention to cultural responsiveness and sustainability.

Mindfulness and psychological health in adults

Mindfulness research in adults provides the foundation for understanding its potential with younger populations. Much of the early empirical

Received: 12 January 2026 | Accepted: 09 March 2026 | Published: 01 June 2026
© The Author(s). Published by Psychreg Ltd under CC BY-NC-ND 4.0.

evidence emerged from clinical psychology and behavioural medicine, where MBIs were developed to alleviate stress and reduce relapse in chronic conditions (Goldberg, 2022).

MBSR, developed by Jon Kabat-Zinn in the late 1970s, was originally designed for patients with chronic pain. It integrates mindfulness meditation, body scans, and gentle yoga across an eight-week group programme. Numerous clinical trials and meta-analyses have since shown that MBSR is associated with significant reductions in stress, anxiety, and depressive symptoms, as well as improvements in quality of life (Grossman et al., 2004; Khoury et al., 2015). These findings helped establish mindfulness as a credible, evidence-based intervention rather than a practice confined to contemplative traditions (Mohamed et al., 2023).

MBCT was developed specifically as a relapse-prevention intervention for recurrent depression. By combining cognitive therapy techniques with mindfulness practice, it helps participants develop non-judgemental awareness of thoughts and feelings, reducing rumination and vulnerability to relapse (Tse & Xu, 2025). Randomised controlled trials have shown that MBCT significantly lowers the risk of depressive relapse among individuals with three or more prior episodes (Kuyken et al., 2016; Segal et al., 2002), underscoring its value in strengthening emotional regulation and interrupting maladaptive cognitive patterns (Bautista et al., 2018).

Research has also extended to non-clinical adult populations. Mindfulness practice consistently shows benefits for psychological well-being, including gains in self-awareness, resilience, and adaptive coping (Creswell, 2017; Relojo-Howell, 2020). Workplace studies have demonstrated reductions in burnout and improvements in concentration and job satisfaction following mindfulness training (Hülsheger et al., 2013). The American Psychological Association (2019) has described mindfulness meditation as a research-supported method of stress reduction, reflecting its broad acceptance within mainstream psychological practice.

At the mechanistic level, mindfulness appears to influence multiple psychological and physiological processes. Cognitively, it reduces habitual rumination and supports greater flexibility in responding to stressors (Keng et al., 2011). Emotionally, it enhances regulation by increasing awareness of internal states and reducing reactivity. Neuroimaging research has identified structural and functional changes in regions associated with attention and emotional processing, including the prefrontal cortex, anterior cingulate cortex, and amygdala (Tang et al., 2015). Physiological markers such as reduced cortisol levels and improved immune functioning further support its biopsychosocial benefits.

That said, limitations in this literature warrant acknowledgement. Many studies rely on self-report measures, raising concerns about reporting bias and placebo effects. Intervention designs vary considerably, complicating cross-study comparisons. Long-term adherence remains an open question: short-term benefits are reasonably well established, but sustaining practice beyond structured programmes is less well documented. These caveats are worth keeping in mind as the field moves to apply mindfulness with younger populations.

Overall, research with adults demonstrates that mindfulness is a flexible, evidence-based approach that can improve mental health and well-being across diverse contexts. This foundational body of work informs efforts to extend mindfulness to younger populations, particularly adolescents, who may benefit from similar emotional and cognitive supports during a formative stage of development.

Mindfulness-based interventions and adolescent mental health

Adolescence is a period of rapid biological, cognitive, and socio-emotional change. It is also a time of heightened risk: the World Health Organization (2021) estimates that approximately one in seven adolescents globally experiences a mental disorder, underscoring the urgent need for effective, accessible interventions. Against this backdrop, MBIs have attracted growing attention as a potentially feasible and scalable approach to supporting youth well-being.

The fit between mindfulness and adolescent development is conceptually plausible. The core tasks of this period, including identity formation, emotional regulation, and managing complex peer dynamics, map closely onto the skills that mindfulness cultivates. Present-moment awareness and non-judgemental acceptance of thoughts and feelings may help buffer against rumination and avoidance; two patterns closely associated with adolescent anxiety and depression. Delivery adaptations such as shorter sessions and more interactive formats are generally considered

necessary to suit the cognitive and motivational characteristics of this age group (Zoogman et al., 2015).

Empirical support, while still developing, is encouraging. Goyal et al. (2023) conducted a pilot study with adolescents and found improvements in attention regulation, stress reduction, and overall psychological functioning. Crucially, the study demonstrated feasibility: participants could engage meaningfully with mindfulness practices when these were appropriately adapted. Jobin et al. (2025) conducted a broader scoping review synthesising evidence across diverse cultural and educational settings, concluding that MBIs show promise in reducing anxiety and depressive symptoms while supporting resilience, self-regulation, and peer relationships. The review also identified significant gaps, particularly the need for larger randomised controlled trials, longer follow-up periods, and culturally sensitive programme designs.

Albright (2023) adds an important dimension by focusing on underserved youth, a population that often faces pronounced barriers to conventional mental health care. Results indicated that MBIs enhanced coping strategies, reduced behavioural difficulties, and produced positive mental health outcomes, suggesting that mindfulness may help address equity gaps in access to psychological support. Keng et al. (2011) provide broader corroborating evidence that mindfulness benefits psychological health across populations, lending additional weight to its application during adolescence. The American Psychological Association (2019) similarly recognises mindfulness meditation as a research-supported method of stress reduction, lending credibility to its use in schools and community programmes for youth.

The mechanisms through which MBIs exert their effects in adolescents are thought to parallel those observed in adults, but with developmental nuances. Mindfulness may reduce stress by decreasing physiological arousal and enhancing parasympathetic nervous system activity. Psychologically, MBIs cultivate attention control, allowing adolescents to disengage from negative thought cycles. Emotionally, mindfulness encourages acceptance, reducing avoidance behaviours and fostering resilience. In group-based settings, mindfulness practice may also promote empathy, prosocial behaviour, and improved peer relationships.

Despite promising results, research on MBIs in adolescents is still developing. Pilot studies often have small sample sizes, limiting generalisability. Variability in intervention design across duration, delivery format, and facilitator expertise complicates comparisons across studies. Engagement remains another challenge, as adolescents may struggle with sustained practice outside structured sessions. Cultural factors may further influence acceptability; mindfulness practices rooted in contemplative traditions may require adaptation to align with the values and experiences of diverse youth populations.

The current evidence base supports cautious optimism, but it does not yet justify strong efficacy claims. Unlike the adult literature, where MBSR and MBCT have been evaluated through large-scale trials with active comparators and long follow-up, adolescent research remains largely preliminary. Small sample sizes, heterogeneous intervention designs, and reliance on self-report measures continue to challenge the strength of conclusions.

These limitations do not, however, reduce the urgency of action. The adolescent mental health crisis requires approaches that are accessible, cost-effective, and developmentally appropriate. Mindfulness fulfils these criteria by offering strategies for attention regulation, stress reduction, and emotional resilience that can be embedded into daily routines. The fact that MBIs have shown benefits even in underserved populations (Albright, 2023) suggests they can help close equity gaps in mental health care if adapted responsibly.

The more productive question at this stage is not whether mindfulness works, but under what conditions it works well enough, for whom, and in what form. Shorter, more interactive sessions tend to fare better than traditional eight-week courses. Technology-enabled delivery through mobile applications or online modules can extend reach, particularly in underserved areas, though the evidence base for digital MBIs in youth remains thin and the risk of low-quality provision is real. School-based integration holds particular promise because it positions mindfulness as a normative developmental skill rather than a clinical remedy, potentially reducing stigma and widening uptake. Cultural adaptation deserves more systematic attention than it has so far received; assumptions about acceptability should not be imported uncritically from Western research contexts into culturally distinct settings. Future research should

treat cultural fit as a design variable rather than an afterthought and should prioritise implementation science over further proof-of-concept work.

This perspective highlights mindfulness as more than a therapeutic technique: it is a developmental resource with the potential to support adolescents in navigating one of the most critical stages of life. Evidence from adults confirms the effectiveness of MBIs in reducing stress, anxiety, and depression, while emerging adolescent research signals comparable benefits in emotional regulation, resilience, and coping.

Yet the value of mindfulness for young people extends beyond symptom relief. By equipping adolescents with present-moment awareness and non-judgemental acceptance, MBIs can lay the foundation for lifelong well-being. The challenge ahead lies in ensuring these interventions are accessible, engaging, and culturally sensitive. Schools, community organisations, and digital platforms are promising avenues for widespread dissemination, but they require thoughtful adaptation.

It is argued that the adolescent mental health crisis demands innovative solutions, and mindfulness represents one such solution ready for broader application. Future work should prioritise scalable models, long-term follow-up, and policy-level support. If approached strategically, mindfulness can move from the margins of clinical practice to the centre of youth development, empowering adolescents not only to manage distress but to thrive.

Acknowledgements: None declared
Conflict of interests: None declared
Ethical approval: Not applicable
Funding: None declared

REFERENCES

Albright, M. G. (2023). Effects of mindfulness-based intervention on mental health outcomes in youth who are underserved [Doctoral dissertation, University of Alabama at Birmingham]. UAB Digital Commons. https://digitalcommons.library.uab.edu/etd-collection/455

American Psychological Association. (2019, October 30). Mindfulness meditation: A research-proven way to reduce stress. https://www.apa.org/topics/mindfulness/meditation

Bautista, L., Relojo, D., Pilao, S.J., Tubon, G., & Andal, M. (2018). Link between lifestyle and self-regulated development as components of academic performance: Basis for a psychoeducational intervention. *Journal on Educational Sciences & Psychology, 8*(52), 68–72. https://doi.org/f8nb

Bear, H. A., Ayala Nunes, L., Ramos, G., Manchanda, T., Fernandes, B., Chabursky, S., Walper, S., Watkins, E., & Fazel, M. (2024). The acceptability, engagement, and feasibility of mental health apps for marginalized and underserved young people: Systematic review and qualitative study. *Journal of Medical Internet Research, 26*, e48964. https://doi.org/10.2196/48964

Creswell, J. D. (2017). Mindfulness interventions. *Annual Review of Psychology*, *68*(2017), 491–516. https://doi.org/10.1146/annurev-psych-042716-051139

Goldberg, S. B. (2022). A common factors perspective on mindfulness-based interventions. *Nature Reviews Psychology*, *1*(10), 605–619. https://doi.org/10.1038/s44159-022-00090-8

Goyal, P., Chandra, M., Rushi, & Choudhary, M. (2023). The effect of mindfulness practice on adolescents: A pilot study. *Journal of Indian Association for Child and Adolescent Mental Health, 19*(2), 207–214. https://doi.org/10.1177/09731342231184628

Grossman, E. D., Blake, R., & Kim, C. Y. (2004). Learning to see biological motion: Brain activity parallels behavior. *Journal of Cognitive Neuroscience*, *16*(9), 1669–1679. https://doi.org/10.1162/0898929042568569

Hülsheger, U. R., Alberts, H. J., Feinholdt, A., & Lang, J. W. (2013). Benefits of mindfulness at work: The role of mindfulness in emotion regulation, emotional exhaustion, and job satisfaction. *Journal of Applied Psychology*, *98*(2), 310–325. https://doi.org/10.1037/a0031313

Jobin, K., Ashok, L., Manjula, M., JJ, T. A., Mathias, E. G., & Krishnan, P. (2025). Mindfulness-based interventions for enhancing adolescent mental health and well-being: A scoping review. *Clinical Epidemiology and Global Health*, *32*, 101961. https://doi.org/10.1016/j.cegh.2025.101961

Kabat-Zinn, J. (1994, September). Catalyzing movement towards a more contemplative/sacred-appreciating/non-dualistic society. In *Meeting of the working group*.

Keng, S. L., Smoski, M. J., & Robins, C. J. (2011). Effects of mindfulness on psychological health: A review of empirical studies. *Clinical Psychology Review, 31*(6), 1041–1056. https://doi.org/10.1016/j.cpr.2011.04.006

Khoury, B., Sharma, M., Rush, S. E., & Fournier, C. (2015). Mindfulness-based stress reduction for healthy individuals: A meta-analysis. *Journal of Psychosomatic Research*, *78*(6), 519–528. https://doi.org/10.1016/j.jpsychores.2015.03.009

Kuyken, W., Warren, F. C., Taylor, R. S., Whalley, B., Crane, C., Bondolfi, G., Hayes, R., Huijbers, M., Ma, H., Schweizer, S., Segal, Z., Speckens, A., & Teasdale, J. D. (2016). Efficacy of mindfulness-based cognitive therapy in prevention of depressive relapse: An individual patient data meta-analysis from randomized trials. *JAMA Psychiatry, 73*(6), 565–574. https://doi.org/10.1001/jamapsychiatry.2016.0076

Mohamed, N. H., Beckstein, A., Winship, G., Ashraf Khan Mou, T., Pang, N. T. P., & Relojo-Howell, D. (2023). Effects of self-expressive writing as a therapeutic method to relieve stress among university students. *Journal of Poetry Therapy*, 1–13. https://doi.org/10.1080/08893675.2023.2174678

Ogwuche, C.H., Caleb, O., & Relojo-Howell, D. (2020). Perceived stress and social support as predictors of subjective well-being among university students in Nigeria. *Psychology & Society, 1*(79), 120–125. https://doi.org/10.35774/pis2020.01.120

Relojo-Howell, D. (2020). Bouncing back, blogging knack: A general overview of resilience and blogging as a psychological intervention. *Central Mindanao University Journal of Science, 24*(2), 5–6, https://doi.org/10.52751/hiqp9266

Segal, Z. V., Williams, J. M. G., & Teasdale, J. D. (2002). Mindfulness-based cognitive therapy for depression: A new approach to preventing relapse. Guilford Press.

Tang, Y. Y., Hölzel, B. K., & Posner, M. I. (2015). The neuroscience of mindfulness meditation. *Nature Reviews Neuroscience, 16*(4), 213–225. https://doi.org/10.1038/nrn3916

Tse, B., & Xu, J. Q. (2025). History and Theoretical Framework of Mindfulness-Based Cognitive Therapy. In *The Palgrave handbook of third-wave psychotherapies* (pp. 455-488). Springer Nature Switzerland. https://doi.org/10.1007/978-3-032-00353-9_20

World Health Organization. (2021). Adolescent mental health. https://www.who.int/news-room/fact-sheets/detail/adolescent-mental-health

Zoogman, S., Goldberg, S. B., Hoyt, W. T., & Miller, L. (2015). Mindfulness interventions with youth: A meta-analysis. *Mindfulness, 6*(2), 290–302. https://doi.org/10.1007/s12671-013-0260-4